

Hubungan penggunaan konsentrasi pil terhadap usia menopause di kota Depok tahun 2012

Fitriyani

Deskripsi Lengkap: <https://lib.fkm.ui.ac.id/detail.jsp?id=91032&lokasi=lokal>

Abstrak

Abstrak
Menopause merupakan menstruasi yang berhenti secara permanen yang<div> disebabkan kehilangan fungsi folikel sel-sel telur. Wanita yang memasuki</div><div> menopause mengalami penurunan hormon estrogen yang mengganggu</div><div> aktivitas sehari-hari, bahkan menurunkan kualitas hidup. Penggunaan kon-</div><div> trasepsi pil berhubungan dengan penundaan usia dan keluhan menopause.</div><div> Penelitian ini bertujuan mengetahui hubungan antara penggunaan kon-</div><div> trasepsi pil terhadap usia menopause. Penelitian ini menggunakan desain</div><div> potong lintang. Populasi adalah wanita menopause di Pos Pembinaan</div><div> Terpadu (Posbindu) Kota Depok. Sampel pada penelitian adalah wanita</div><div> menopause yang berusia 45 _ 60 tahun. Teknik pengambilan sampel se-</div><div> cara purposive sampling subjek dengan besar sampel 407 orang. Analisis</div><div> multivariat pada penelitian ini menggunakan cox proportional hazard model.</div><div> Hasil analisis multivariat menunjukkan tidak ada hubungan antara lama</div><div> penggunaan kontrasepsi pil terhadap usia menopause baik sebelum mau-</div><div> pun sesudah dikontrol variabel kovariat, yaitu tingkat pendidikan. Namun</div><div> demikian, masih diperlukan penelitian lain dengan menggunakan desain</div><div> penelitian kohort prospektif untuk dapat melihat hubungan temporal antara</div><div> lama penggunaan kontrasepsi pil terhadap usia menopause.</div><div> Menopause is marked with the permanent cessation of menstruation due to</div><div> the loss of follicles. Earlier menopause will be likely to increase the risk</div><div> factors relating to declined estrogen level, such as osteoporosis that can</div><div> lead to early death. A woman entering menopause period often experiences</div><div> declined estrogen hormone that causes her to have complaints or distur-</div><div> ances that hinder her daily activities and even reduce her quality of life.</div><div> However, the use of oral contraceptive poses a correlation with the post-</div><div> poning of menopause age and complaints. The primary aim of this study</div><div> was to examine the relation of oral contraceptive use and age at</div><div> menopause. This was an observational study with cross-sectional study</div><div> design. Population in this study was all menopausal women in integrated</div><div> training post (Posbindu), Depok. The sample was menopausal women</div><div> </div><div> </div>among 45 _ 60 years old. Sample was 407 menopausal women taken<div> purposive sampling. The data was analysed by cox?s proportional hazard</div><div> analysed. The longer use of oral contraceptive was not associated with age</div><div> at menopause before and after adjusted for confounding variable (educa-

tion). However, another similar studies was still needed with prospective
cohort study design to know temporality causal of longer use of
oral contraceptive and age at menopause.